

ON V-SYMMETRY IN SU (3)

K. D. KRORI

DEPARTMENT OF PHYSICS,
COTTON COLLEGE, GAUHATI, ASSAM, INDIA

(Received April 26, 1966; Resubmitted June 6, 1968)

ABSTRACT. Some properties of V -symmetry in $SU(3)$ have been discussed here.

INTRODUCTION

V -symmetry is one of the subgroups of $SU(3)$. It appears that in different pieces of literature (Dalitz, 1963; Mayer, 1963; Low, 1964; Sakurai, 1964; Lipkin, 1966; Matthews, 1966; Swart, 1966), the other two subgroups, I -symmetry and U -symmetry, have been studied in comparatively greater detail. We propose to consider certain aspects of V -symmetry in this paper.

Now we briefly introduce V -symmetry in a general way. Let us consider three unit vectors in F_3 — F_8 or I_3 — Y plane [F_i being the infinitesimal generators in the basic representation of $SU(3)$]:

$$\vec{i} = (1, 0), \quad \vec{u} = (-\frac{1}{2}, \sqrt{3}/2), \quad \vec{v} = (\frac{1}{2}, \sqrt{3}/2) \quad \dots (1)$$

and define the vector generator :

$$\vec{E} = (F_3, F_8) \quad \dots (2)$$

and put

$$\vec{i} \cdot \vec{E} = I_3, \quad \vec{u} \cdot \vec{E} = U_3, \quad \vec{v} \cdot \vec{E} = V_3 \quad \dots (3)$$

Then,

$$I_3 + U_3 - V_3 = 0 \quad \dots (4)$$

Also we define :

$$\begin{aligned} I_{\pm} &= F_1 \pm iF_2 \\ U_{\pm} &= F_6 \pm iF_7 \\ V_{\pm} &= F_4 \pm iF_5 \end{aligned} \quad \dots (5)$$

Thus we obtain the three subgroups (I , I_3), (U , U_3) and (V , V_3), of which the first is just iso-spin (I -spin) and the others are U -spin and V -spin respectively. (I_1 , I_2), (U_1 , U_2) and (V_1 , V_2) are given by :

$$\begin{aligned} I_1 &= \frac{I_+ + I_-}{2} & I_2 &= \frac{I_+ - I_-}{2} & U_1 &= \frac{U_+ + U_-}{2} & U_2 &= \frac{U_+ - U_-}{2} \\ V_1 &= \frac{V_+ + V_-}{2} & V_2 &= \frac{V_+ - V_-}{2} \end{aligned} \quad (6)$$

PARTICLES AT ORIGIN OF WEIGHT DIAGRAMS

In a weight diagram (figure 1), just as the horizontal lines parallel to \vec{i} link particles of the same I -spin multiplets, so the lines parallel to \vec{v} connect V -multiplets. Examples of V -spin multiplets are :

$$(K^0, \pi^-), (\pi^+, \bar{K}^0), (n, \Sigma^-), (\Sigma^+, \Xi^0) \text{—(fig. 2 and 5)}$$

$$(\Delta^{++}, Y_1^{*+}, \Xi^{*0}, \Omega^-)(Y^{*++}, \Xi^{*+0}) \text{—(fig. 3 and 4).}$$




Figure 1,




Figure 2.




Figure 3.




Figure 4,

Where there are two particles at the origin on a weight diagram (say, figure 5), we may define in the F_3-F_8 plane a vector (Matthews)

$$\vec{\pi^0} = (\pi^0, \eta)$$

Then, the $I_3 = 0$ component of the I -spin triplet is

$$\pi^0_i = \vec{\pi^0} \cdot \vec{i} = \pi^0,$$




Fig. 5.

and the I -spin singlet orthogonal to it is

$$\eta_t = \eta$$

Similarly, the $V_3 = 0$ component of the V -spin triplet (figure 5) is from (1) :

$$\pi_v^0 = \vec{\pi}^0 \cdot \vec{v} = \frac{\pi^0 + \sqrt{3}}{2} \quad \dots (7)$$

and

$$\eta_v = \frac{\sqrt{3}\pi^0 + \eta}{2}$$

Also, for the baryon octet (figure 2)

$$\Sigma_v^0 = \frac{\Sigma^0 + \sqrt{3}\Lambda}{2}$$

$$\Lambda_v = \frac{\sqrt{3}\Sigma^0 + \Lambda}{2} \quad \dots (8)$$

and for the baryon resonance (figure 4)

$$Y_{1v}^{*0} = \frac{Y_1^{*0} + \sqrt{3}Y}{2}$$

$$Y_{0v}^{*0} = \frac{\sqrt{3}Y_1^{*0} + Y_0^{*0}}{2} \quad \dots (9)$$

V_2 -SYMMETRY

Following the cases of I -spin and U -spin, let us consider the discrete transformation :

$$P_v = \exp[i V_2 \pi] \quad \dots (10)$$

This will have the effect of reflecting the weight diagrams in the $V_3 = 0$ axis. Then, we should have the following transformations :

$$\begin{aligned} K^0 & \leftrightarrow \pi^-, K^+ \leftrightarrow K^-, \pi^+ \leftrightarrow \bar{K}^0, n \leftrightarrow \Sigma^-, p \leftrightarrow \Xi^-, \Sigma^+ \leftrightarrow \Xi^0, \\ \Delta^0 & \leftrightarrow Y_1^{*-}, \Delta^+ \leftrightarrow \Xi^-, \Delta^{++} \leftrightarrow \Omega^- \end{aligned}$$

Invariance under these transformations implies that

$$\langle f | S | i \rangle = \langle f | P_v^\dagger S P_v | i \rangle \quad \dots (11)$$

where $\langle f | S | i \rangle$ stands for the S -matrix between the initial state $|i\rangle$ and the final state $\langle f|$. Then we should expect :

$$\begin{aligned} \langle \pi^- \Sigma^+ | S | \Sigma^- \pi^+ \rangle &= \langle K^0 \Xi^0 | S | n \bar{K}^0 \rangle \\ \langle K^- p | S | \Xi^- K^+ \rangle &= \langle K^+ \Xi^- | S | p K^- \rangle \end{aligned} \quad \dots (12)$$

Again, since $\Lambda_p \leftrightarrow \Lambda_n$ under P_v -transformation, we may take $\Lambda \leftrightarrow \Lambda$ from (8). This should give :

$$\begin{aligned} \langle \Lambda K^0 | S | p \pi^- \rangle &= \langle \Lambda \pi^- | S | \Xi^- K^0 \rangle \\ \langle \Lambda K^+ | S | n \pi^+ \rangle &= \langle \Lambda K^- | S | \Sigma^- \bar{K}^0 \rangle \end{aligned} \quad \dots (13)$$

MAGNETIC MOMENTS

For any V -multiplet, the charge is given by :

$$Q = V_3 + a \quad \dots (14)$$

Thus, for the doublet (Σ^+, Ξ^0) , $a = \frac{1}{2}$. For the triplet (p, Σ_v^0, Ξ^-) , $a = 0$. For the quadruplet $(\Delta^{++}, Y_1^{*+}, \Xi^{*0}, \Omega^-)$ also, $a = \frac{1}{2}$.

Now, since Q depends linearly upon V_3 , any physical property dependent linearly on electromagnetic interaction may be expressed in the form :

$$\alpha + \beta V_3 \quad \dots (15)$$

for a V -spin multiplet. The static magnetic moments for the triplet (p, Σ_v^0, Ξ^-) will be given by

$$\mu(p) - \mu(\Sigma_v^0) = \mu(\Sigma_v^0) - \mu(\Xi^-) \quad \dots (16)$$

Now, taking (Mayer)

$$\mu(\Xi^-) = -\mu(p) - \mu(n)$$

we have from (16)

$$\mu(\Sigma_v^0) = -\frac{1}{2}\mu(n) \quad \dots (16a)$$

On the other hand (Matthews)

$$\mu(\Sigma_u^0) = \mu(n) = -2\mu(\Sigma_v^0)$$

The full expression for $\mu(\Sigma_v^0)$ may be written thus

$$\begin{aligned} \mu(\Sigma_v^0) &= \langle \Sigma_v^0 | J | \Sigma_v^0 \rangle = \left\langle \frac{\Sigma^0 + \sqrt{3}\Lambda}{2} \left| J \right| \frac{\Sigma^0 + \sqrt{3}\Lambda}{2} \right\rangle \\ &= \frac{1}{4}\mu(\Sigma^0) + \frac{3}{4}\mu(\Lambda) + \frac{\sqrt{3}}{2}\mu(\Lambda\Sigma^0) \end{aligned} \quad (17)$$

Now, it can be shown (Matthews) that

$$\frac{\sqrt{3}}{2}\mu(\Lambda\Sigma^0) = \frac{3}{4}\mu(\Sigma^0) - \frac{3}{4}\mu(\Lambda) \quad \dots (18)$$

combining (17) and (18),

$$\mu(\Sigma_v^0) = \mu(\Sigma^0) \quad \dots (19)$$

Also, for the triplet (Δ^+ , Y_1^{*0} , Ξ^{*-})

$$\begin{aligned} &\mu(\Delta^+) - \mu(Y_1^{*0}) \\ &= \mu(Y_1^{*0}) - \mu(\Xi^{*-}) \end{aligned} \quad \dots (20)$$

and for the quadruplet (Δ^{++} , Y_1^{*+} , Ξ^{*0} , Ω^-)

$$\mu(\Delta^{++}) - \mu(Y_1^{*+}) = \mu(Y_1^{*+}) - \mu(\Xi^{*0}) = \mu(\Xi^{*0}) - \mu(\Omega^-) \quad \dots (21)$$

We propose to obtain with V -symmetry some well-known mass-relations which have already been derived otherwise.

(a) *Parallelogram and hexagon laws*

MASS RELATIONS

First, we consider parallelogram law with reference to I -spin and V -spin. If there is only one particle at each point in the weight diagram (fig.-1), the law gives :

$$m(1) - m(2) + m(3) - m(4) = 0 \quad \dots (22)$$

This follows I -spin conservation

$$m(1) = m(2), \quad m(3) = m(4) \quad \dots (23)$$

and *V*-spin conservation

$$m(1) = m(4), \quad m(2) = m(3) \quad (24)$$

For the baryon decuplet, we have according to (22)

$$\begin{aligned} \Delta^+ - \Delta^{++} + Y_1^{*0} - Y_1^{*-} &= 0 \\ Y_1^{*0} - Y_1^{*+} + \Xi^{*0} - \Xi^{*-} &= 0 \quad \dots \quad (25) \\ \Delta^0 - \Delta^+ + Y_1^{*-} - Y_1^{*0} &= 0 \end{aligned}$$

If, however, there are two particles, say at the point, they may be denoted in *I*-space by and (5) and in *V*-space by and (5_v). Now, we have the relations :

$$\begin{aligned} m(1) - m(2) + m - m(4) + \alpha m(5) \\ m(1) - m(2) + \beta m(3_v) - m(4) + \gamma m(3_v, 5_v) = 0 \end{aligned} \quad (26)$$

where $m(35)$ and $m(3_v, 5_v)$ are transition masses in the *I* and *V* representations respectively. The constants may be found from the identity (26). However, for *V*-spin conservation only, β is assumed to be = 1 to satisfy (24).

For the baryon octet, there are two parallelograms. Let us consider the parallelogram (*n*, *p*, Σ^0 , Λ , Σ^-).

From (26)

$$\begin{aligned} n - p + \Sigma^0 - \Sigma^- + \alpha(\Sigma^0 \Lambda) \\ n - p + \beta \Sigma^0 - \Sigma^- + \gamma(\Sigma^0 \Lambda_v) = 0 \end{aligned} \quad \dots \quad (27)$$

It is found from (8) and (27) that (Appendix)

$$\alpha = \sqrt{3}, \quad \beta = 2 \text{ and } \gamma = 2\sqrt{3} \quad \dots \quad (23)$$

[It may be stated in passing that for the *I* - *U* parallelogram (*n*, *p*, Σ^+ , Σ^0), the values are found to be (Appendix-) $\alpha = \sqrt{3}$, $\beta = 1$, $\gamma = \sqrt{3}$.]

Thus,

$$n - p + \Sigma^0 - \Sigma^- + \sqrt{3}(\Sigma^0 \Lambda) = 0 \quad \dots \quad (29)$$

Similarly, for the other parallelogram.

$$\Sigma^0 - \Sigma^+ + \Xi^0 - \Xi^- + \sqrt{3}(\Sigma^0 \Lambda) = 0 \quad (30)$$

(29) and (30) quite satisfactorily agree with the corresponding relations obtained with *I* - *U* parallelogram (Matthews). From (29) and (30), we obtain the well-known six-mass relation :

$$n - p + \Sigma^+ - \Sigma^- + \Xi^- - \Xi^0 = 0 \quad \dots \quad (31)$$

This relation may also be directly obtained by an extension of parallelogram to hexagon considering conservation of I -spin, U -spin and V -spin. Thus we have (figures 2, 4, 5) :

$$m(1) - m(2) + m(3) - m(4) + m(5) - m(6) = 0$$

This applies to baryon, meson and baryon resonance octets. It thus appears that the six-mass relation is a characteristic of an octet.

(b) *Electromagnetic and medium-strong interaction effects*

Like I -symmetry, V -symmetry is broken by electromagnetic interactions, whereas like U -symmetry, it is violated by medium-strong interactions. In analogy with other case, it is plausible to write (Matthews) the mass formula as—

$$m = \alpha + \beta V_3 + \gamma V_3^2 \quad (32)$$

Then, for the quadruplet (Δ^{++} , Y_1^{*+} , Ξ^{*0} , Ω^-)

$$\Delta^{++} - 2Y_1^{*+} + \Xi^{*0} = Y_1^{*+} - 2\Xi^{*0} + \Omega^- \quad (33)$$

Now,

$$\left. \begin{aligned} \text{L.H.S.} &= 1236 - 2 \times 1382.7 + 1529.7 = +0.3 \text{ Mev} \\ \text{R.H.S.} &= 1382.7 - 2 \times 1529.7 + 1675 = -1.7 \text{ Mev} \end{aligned} \right\} \text{ (Swart)}$$

The discrepancy between the two sides is to the extent of 2Mev, i.e., about 0.13% only of the average mass.

Following Okubo (Matthews), if it is assumed in this case also that the first-order term dominates in (32), then we have

$$m = \alpha + \beta V_3 \quad \dots (34)$$

Applying (34) to the above quadruplet, we obtain exactly the equal spacing rule (Swart) :

$$\Delta^{++} - Y_1^{*+} = Y_1^{*+} - \Xi^{*0} = \Xi^{*0} - \Omega^- \quad \dots (35)$$

That is,

$$\left. \begin{aligned} \Delta^{++} - Y_1^{*+} &= 1236 - 1382.7 = -146.7 \\ Y_1^{*+} - \Xi^{*0} &= 1382.7 - 1529.7 = -147.0 \\ \Xi^{*0} - \Omega^- &= 1529.7 - 1675 = -145.3 \end{aligned} \right\} \text{ (Swart)}$$

We consider the baryon triplet (p , Σ_p^0 , Ξ^-) in the light of (34) and obtain

$$p - \Sigma_p^0 = \Sigma_p^0 - \Xi^- \quad \dots (36)$$

Now,

$$\Sigma_v^0 = \frac{1}{4} \Sigma^0 + \frac{3}{4} \Lambda + \frac{\sqrt{3}}{5} (\Sigma^0 \Lambda) \quad \dots (37)$$

(from Appendix)

From (36) and (37)

$$\frac{1}{2}(p + \Xi^-) = \Sigma_v = \frac{1}{4} \Sigma^0 + \frac{3}{4} \Lambda + \frac{\sqrt{3}}{5} (\Sigma^0 \Lambda) \quad \dots (38)$$

Since the average value of the transition mass ($\Sigma^0 \Lambda$) is about 1.15 Mev from (29) and (30), it may be neglected compared to other masses and we obtain

$$\frac{1}{2}(p + \Xi^-) = \frac{1}{4} \Sigma^0 + \frac{3}{4} \Lambda \quad \dots (39)$$

This is just a form of the well-known baryon mass-relation (Swart, Mayor) :

$$\frac{1}{2}(M_N + M_\Xi) = \frac{1}{4}(M_\Sigma + 3M_\Lambda) \quad \dots (40)$$

Applying (34) to the meson triplet (K^+ , π_v^0 , K^-) we have as above (masses squared)

$$\frac{1}{2}(K^+ + K^-) = \pi_v^0 = \frac{1}{4} \pi^0 + \frac{3}{4} \eta + \frac{\sqrt{3}}{2} (\pi^0 \eta) \quad \dots (41)$$

Neglecting the transition mass ($\pi^0 \eta$) as before,

$$\frac{1}{2}(K^+ + K^-) = \frac{1}{4} \pi^0 + \frac{3}{4} \eta \quad \dots (42)$$

This is also a form of the well-known meson mass-relation (Swart, Mayer) :

$$m_K^2 = \frac{1}{4}(m_\pi^2 + 3m_\eta^2) \quad \dots (42)$$

It is interesting that a number of mass-relations may be obtained from simple premises on *V*-symmetry. The above results also vindicate Okubo's suggestion which has been extended to *V*-spin here.

ELECTROMAGNETIC INTERACTIONS

Let us take the following interactions :

$$\begin{aligned} \Sigma^0 &\rightarrow \pi + \gamma \\ \eta &\rightarrow \pi^+ + \pi^- + \gamma \\ \gamma + p &\rightarrow n + \pi^+ \\ \gamma + p &\rightarrow \pi + K^+ \end{aligned} \quad \dots (43)$$

It appears that in these processes, $|\Delta V_3| = 0$. Thus, V_3 is conserved in electromagnetic interactions.

WEAK INTERACTIONS

Let us consider the weak interactions :

$$\Delta \rightarrow p + \pi^-, \quad n + \pi^0$$

$$\Sigma^+ \rightarrow p + \pi^0,$$

$$K \rightarrow 2\pi, 3\pi \text{ etc.}$$

$$\Sigma^- \rightarrow n + \pi^-$$

$$\Xi^- \rightarrow \Lambda + \pi^- \quad (44)$$

and so on

In all cases, the change in V_3 appears, with reference to (7) and (8), to be

$$|\Delta V_3| = \frac{1}{2} \quad \dots (45)$$

It appears that V -spin is also similar to I -spin in respect of weak interactions. This is quite in agreement with (4).

ACKNOWLEDGEMENTS

The author expresses his profound gratitude to the Government of Assam for all facilities provided at Cotton College, Gauhati and to the University Grants Commission, Delhi, for all sorts of encouragement in working out this paper.

APPENDIX

From (8), we have for mass,

$$\Sigma_v^0 = \langle \Sigma_v^0 | m | \Sigma_v^0 \rangle = \frac{1}{4} \Sigma^0 + \frac{3}{4} \Lambda + \sqrt{3} (\Sigma^0 \Lambda)$$

$$(\Sigma_v^0 \Lambda_v) = \langle \Sigma_v^0 | m | \Lambda_v \rangle = \frac{\sqrt{3}}{4} \Sigma^0 + \frac{\sqrt{3}}{4} \Lambda + (\Sigma^0 \Lambda)$$

Now, from (30), we have the identity :

$$\begin{aligned} & n - p + \Sigma^0 - \Sigma^- + \alpha (\Sigma^0 \Lambda) \\ & \equiv n - p + \left(\frac{\beta}{4} + \frac{\sqrt{3}}{4} \gamma \right) \Sigma^0 + \left(\frac{3\beta}{4} + \frac{\sqrt{3}}{4} \gamma \right) \Lambda - \Sigma^- + \left(\frac{\sqrt{3}}{2} \beta + \gamma \right) (\Sigma^0 \Lambda) \end{aligned}$$

Hence, we obtain the following equations :

$$\frac{\beta}{4} + \frac{\sqrt{3}}{4} \gamma = 1$$

$$\frac{3\beta}{4} + \frac{\sqrt{3}}{4} \gamma = 0$$

$$\frac{\sqrt{3}}{2} \beta + \gamma = \alpha$$

Solving these equations

$$\alpha = \sqrt{3}, \quad \beta = -2, \quad \gamma = 2\sqrt{3}$$

Also, solving in the same manner the identity for the *I-U* parallelogram (*n*, *p*, Σ^+ , Σ^0), the values obtained are

$$\alpha = \sqrt{3}, \quad \beta = 1, \quad \gamma = \sqrt{3}$$

REFERENCES

- Dalitz, R. H., 1963, *Lectures on the Properties and the Symmetry Theories of the Resonant States*, Tata Institute of Fundamental Research, Bombay, pp. 62-89.
- Lipkin, H. J., 1966, *Lie Groups for Pedestrians*, North-Holland Co., Pp. 143-46.
- Low, F. E., 1964, *Lectures On Elementary Particles and Scattering Theory*, Tata Institute of Fundamental Research, Bombay, Pp. 26-52.
- Mathews, P. T., 1966 (To be Published). Chapter VII-Unitary Symmetry, Group Theory.
- Mayer, M. E., 1963, *Lectures On Strong and Electromagnetic Interactions*, Brandeis Summer Institute in Theoretical Physics, Pp. 263-331.
- Sakurai, J. J., 1964, *Lectures on New Mesons, Resonances and Unitary Symmetry*, Tata Institute of Fundamental Research, Bombay, Pp 69-119.
- Swart, J. J. de, 1966, *Symmetries of Strong Interactions*, CERN.